



SPATHODUS ERYTHRODON KABENZI

Spathodus erythrodon



One of a number of species known as 'Goby Cichlids', these fish hop over the substrate like Gobies. They occur in shallow rocky coastal areas of the lake, a habitat influenced by strong currents from constant wave surges. In the wild they feed on algae and small invertebrates. Provide plenty of aeration and strong filtration, and plenty of rocky habitat.

Disclaimer: The information provided here is of a general nature only. There are site specific factors that are not addressed here that may also affect the health and well being of your fish, as such Coburg Aquarium will not be held liable for the performance of your fish.

Copyright © 2026 Coburg Aquarium

KEEPING SNAPSHOT

Difficulty: Intermediate
Max Growth Size: 8cm
Temperament: Can be territorial

PREFERRED WATER PARAMETERS

Temperature: 23-27 degrees celsius
pH Level: 7.8-9.0
General Hardness: Hard

RANGE

Lake Tanganyika: shallow water surf zone

REPRODUCTION

Mouthbrooder (both male and female may brood with eggs).

FEEDING

Carnivorous and herbivorous. Eats all normal foods.

ADDITIONAL COMMENTS



03 9354 5843



03 9354 9371



info@coburgaquarium.com.au



coburgaquarium.com.au